

NONOBLATENESS OF A GENERATING CONE IN *SH*-SPACE AND ITS APPLICATION

**E. I. SMIRNOV, G. E. KOZLOV, V. V. BOGUN
and A. D. UVAROV**

South Mathematical Institute Russian
Academic of Science and NO-Alania
Yaroslavi State Pedagogical University
150000 Yaroslavi
Respublianskaya 108
Russia
e-mail: smirnov@yspu.org

Abstract

The concept of a nonoblate cone in a Banach space is one of the most important ideas in the theory of ordered normed linear spaces. In connection with the introduction, the new class of *SH*-spaces by Smirnov (the *H*-spaces as Souslin spaces earlier), the problem of clarifying the role of the concept of nonoblateness of a cone in such spaces arises naturally. In the present paper, we will obtain a theorem about the nonoblateness of a generating cone in an *SH*-space and demonstrate a series of its applications to questions of differentiability with respect to a cone and of the continuity of a positive operator. This will allow us to obtain a theorem on the existence of a saddle point of the Lagrange function for linear optimization problems in *SH*-spaces.

2010 Mathematics Subject Classification: 47B65, 46A40, 46A13.

Keywords and phrases: nonoblate cones, locally convex spaces, Kuhn-Tacker theorem, compact differentiability, closed graph theorem.

Communicated by Jong Kwang Yoo.

Received June 27, 2014

1. Nonoblate Cones in *SH*-Spaces

We recall [2] that a cone K in a locally convex space (LCS) X is said to be *nonoblate* if for each neighbourhood of zero U , there exists a neighbourhood of zero V for which $V \subset U \cap K - U \cap K$. The theory of differentiation in an LCS as developed in [1] is used systematically. All topological vector spaces considered are assumed to be separated and locally convex.

Let (G, τ) be a locally convex metric topological vector group (TVG) and K be a closed generating cone in G . We will denote by d a quasinorm defining the topology τ , i.e., a nonnegative functional on G , which satisfies the conditions:

- (a) $0 \leq d(x) \leq 1 \quad (x \in G)$;
- (b) $d(\lambda x) \leq d(x) \quad (|\lambda| \leq 1, x \in G)$;
- (c) $d(x_1 + x_2) \leq d(x_1) + d(x_2) \quad (x_1, x_2 \in G)$.

The quasinorm

$$\tilde{d}(x) = \inf\{d(u) + d(v) : x = u - v, u, v \in K\},$$

defines on G the topology $\tilde{\tau}$ of a locally convex TVG in which a base of absolutely convex neighbourhoods of zero is formed by the sets

$$V_n = K \cap U_n - K \cap U_n \quad (n = 1, 2, \dots),$$

where $\{U_n : n = 1, 2, \dots\}$ is a base of absolutely convex neighbourhoods of zero in the topology τ . It is clear that $\tau \leq \tilde{\tau}$.

Proposition 1. *If (G, τ) is a complete TVG, then it follows from convergence in (G, τ) of the series:*

$$x = \sum_{n=1}^{\infty} x_n, \tag{1}$$

that

$$\tilde{d}(x) \leq \sum_{n=1}^{\infty} \tilde{d}(x_n). \quad (2)$$

Proof. Suppose that the series (1) converges in (G, τ) and the right-hand side of inequality (2) is finite. Then for every $\epsilon > 0$, there exist sequences $u_n \in K$ and $v_n \in K$ for which $x_n = u_n - v_n$ and

$$d(u_n) + d(v_n) \leq \tilde{d}(x_n) + 2^{-n} \epsilon.$$

Since (G, τ) is a complete TVG, it follows from this that there exist elements $u, v \in K$ for which $x = u - v$ and

$$\tilde{d}(x) \leq d(u) + d(v) \leq \sum_{n=1}^{\infty} [d(u_n) + d(v_n)] \leq \sum_{n=1}^{\infty} \tilde{d}(x_n) + \epsilon.$$

Inequality (2) follows from this since $\epsilon > 0$ is arbitrary. The proposition is proved.

From Proposition 1 and the completeness of (G, τ) , we deduce that any series (1) for which the right-hand side of inequality (2) is finite converges in (G, τ) . Hence we have

Proposition 2. *The TVG $(G, \tilde{\tau})$ is complete.*

Proof. Let (x_n) be a fundamental sequence in $(G, \tilde{\tau})$. We choose a subsequence (x_{n_k}) such that

$$\tilde{d}(x_{n_{k+1}} - x_{n_k}) < 2^{-k} \quad (k = 1, 2, \dots).$$

Then

$$\sum_{k=1}^{\infty} \tilde{d}(x_{n_{k+1}} - x_{n_k}) < \infty,$$

and consequently, the series

$$x_{n_1} + \sum_{k=1}^{\infty} (x_{n_{k+1}} - x_{n_k}),$$

converges in $(G, \tilde{\tau})$. In the other words, the subsequence (x_{n_k}) , and along with it also the sequence (x_n) , converge in $(G, \tilde{\tau})$. The proposition is proved.

Theorem 1. *Let (X, τ^*) be an SH-space and K be a generating closed cone in X . Then K is a nonoblate cone.*

Proof. Let (X, τ^*) be an SH-space and K be a generating closed cone in X . Let

$$X = \bigcup_{\nu \in \mathcal{P}} \bigcap_{k=1}^{\infty} X_{n_1 n_2 \dots n_k},$$

be such that τ^* is the strongest locally convex topology on X for which all the embeddings of the locally convex metric TVGs $X_{(\nu)}$ ($\nu \in \mathcal{P}$) in the space (X, τ^*) are continuous. Without loss of generality, it can be assumed that the spaces $X_{n_1 n_2 \dots n_k}$ ($n_k, k = 1, 2, \dots$) are seminormed and the embeddings

$$X_{n_1 n_2 \dots n_{k+1}} \rightarrow X_{n_1 n_2 \dots n_k} \quad (k = 1, 2, \dots; \nu \in \mathcal{P}),$$

are continuous. Here \mathcal{P} is a subset of \mathcal{N}^* , the set of sequences of positive integers.

Let $\nu = (n_1, n_2, \dots) \in \mathcal{P}$. We will denote by $\{U_{n_1 n_2 \dots n_k} : k = 1, 2, \dots\}$ the family of absolutely convex neighbourhoods of zero in a base for the space $X_{(\nu)}$, which are such that for all $k = 1, 2, \dots$, the set $U_{n_1 n_2 \dots n_k}$ is a neighbourhood of zero in $X_{n_1 n_2 \dots n_k}$ and $2U_{n_1 n_2 \dots n_{k+1}} \subset U_{n_1 n_2 \dots n_k}$. Then the sets

$$V_{n_1 n_2 \dots n_k} = U_{n_1 n_2 \dots n_k} \cap K - U_{n_1 n_2 \dots n_k} \cap K \quad (k = 1, 2, \dots),$$

are absolutely convex and their linear hulls $L(V_{n_1 n_2 \dots n_k}) = Y_{n_1 n_2 \dots n_k}$ can be given seminorm topologies in such a way that for each $k = 1, 2, \dots$, the sets ${}^\epsilon V_{n_1 n_2 \dots n_k}$ ($\epsilon > 0$) form a base of neighbourhoods of zero. It is not difficult to see that

$$X = \bigcup_{\nu \in \mathcal{P}} \bigcap_{k=1}^{\infty} Y_{n_1 n_2 \dots n_k},$$

and moreover, the sequence $V_{n_1 n_2 \dots n_k}$ forms a base of absolutely convex neighbourhoods of zero for some TVG $Y_{(\nu)}$. Since the space $X_{(\nu)}$ is complete, then by Proposition 2, the TVG $Y_{(\nu)}$ is also complete.

Now, let us consider on X the strongest locally convex topology σ^* for which all the embeddings of the spaces $Y_{(\nu)}$ ($\nu \in \mathcal{P}$) in the space (X, σ^*) are continuous. Then (X, σ^*) is an *SH*-space and moreover $\tau^* \leq \sigma^*$. By the Closed Graph Theorem for *SH*-spaces, we have the inequality $\sigma^* \leq \tau^*$. The assertion of the theorem now follows since by construction the cone K is nonoblate in (X, σ^*) . The theorem is proved.

Corollary 1. *Let K be a generating closed cone in a sequentially complete bornological *SH*-space (X, τ) . Then K is a nonoblate cone.*

This assertion follows from Theorem 1 and Proposition 7.3.5 of [4].

2. Compact Differentiability with Respect to a Cone

Let X and Y be LCSs, K be a closed cone in X and $L(X, Y)$ be the vector space of all continuous linear mappings from X to Y . We will denote by β (resp., β_c) the system of all bounded (resp., compact) subsets

of the space X , and by β_k (resp., β_c^k) the system of all bounded (resp., compact) subsets of the cone K . Let $L_{\beta}(X, Y)$ (resp., $L_{\beta_c}(X, Y)$) be the LCS obtained by giving the space $L(X, Y)$ the topology of uniform convergence on the sets of the system β (resp., β_c).

We will say (see also [2]) that the operator $A : X \rightarrow Y$ is *differentiable at the point* $x_0 \in X$ *in the directions of the cone* K , if the function $y(t) = A(x_0 + th)$ is differentiable with respect to t at the point $t = 0$ for all $h \in K$. If the derivative $y'(0)$ is representable in the form $y'(0) = A'(x_0)h$ ($h \in K$), where $A'(x_0) \in L(X, Y)$, then we will call the linear operator $A'(x_0)$ the *weak derivative with respect to the cone* K *at the point* x_0 .

If the identity

$$\lim_{t \rightarrow 0} \frac{y(t) - y(0)}{t} = A'(x_0)h,$$

is satisfied uniformly with respect to $h \in B$ for each B from β_k (resp., β_c^k), then we will call $A'(x_0)$ the *bounded* (resp., *compact*) *derivative with respect to the cone* K *at the point* x_0 . Mappings which have a weak, bounded or compact derivative with respect to a cone will be called weakly, boundedly or compactly differentiable with respect to the cone.

Let (X, τ) be a separated sequentially complete bornological *SH*-space, i.e.,

$$X = \bigcup_{\nu \in \mathcal{P}} \bigcap_{k=1}^{\infty} X_{n_1 n_2 \dots n_k},$$

and each space $X_{(\nu)}$ ($\nu \in \mathcal{P}$) is a locally convex complete metric TVG, which is continuously embedded in (X, τ) . The topology τ of the space X induces on each space

$$X_\nu = \bigcap_{k=1}^{\infty} X_{n_1 n_2 \dots n_k},$$

a locally convex topology $\tilde{\tau}_\nu$ which in general is different from the topology τ_ν of the Fréchet space X_ν ($\nu \in \mathcal{P}$). We will assume that $\tau_\nu = \tilde{\tau}_\nu$ for each $\nu \in \mathcal{P}$.

Theorem 2. *Suppose that for the operator $A : X \rightarrow Y$ the weak derivative $A'(x)$ with respect to a generating closed cone K is a continuous mapping into $L_{\beta_c}(X, Y)$ on an open neighbourhood U of the point x . Then $A'(x)$ is the compact derivative of the operator A at the points $x \in U$.*

Proof. By Corollary 1, the cone K is nonoblate in the space (X, τ) and we have the identity

$$X = \bigcup_{\nu \in \mathcal{P}} \bigcap_{k=1}^{\infty} Y_{n_1 n_2 \dots n_k},$$

where the $Y_{n_1 n_2 \dots n_k}$ ($n_k, k = 1, 2, \dots$) are seminormed spaces and the cone K is nonoblate in each locally convex TVG $Y_{(\nu)}$ ($\nu \in \mathcal{P}$). We have to show that

$$\lim_{\delta \rightarrow 0} \frac{A(x + \delta h) - A(x)}{\delta} = A'(x)h, \tag{3}$$

where $x \in U$ and convergence is uniform with respect to all $h \in B$ for every $B \in \beta_c$.

Let $x \in U$, $B \in \beta_c$ and let W be a convex neighbourhood of zero in the space Y . Since the space (X, τ) is sequentially complete, the set B is contained and bounded in some space Y_ν , where $\nu \in \mathcal{P}$. By the Closed Graph Theorem, there exists $\nu' \in \mathcal{P}$ such that $Y_\nu \subset Y_{\nu'}$. But $\tau_\nu = \tilde{\tau}_\nu$; therefore, the set B is compact in Y_ν and thus it is compact in $Y_{\nu'}$. By

Corollary 1 of [3], there is a sequence (h_n) converging to zero in $Y_{\nu'}$ such that B is contained in the closed absolutely convex hull of (h_n) . Because of the nonoblateness of the cone K in the space $Y_{(\nu')}$, there exist sequences $(u_n) \subset K$ and $(v_n) \subset K$ for which $h_n = u_n - v_n$ and $u_n \rightarrow 0$, $v_n \rightarrow 0$ as $n \rightarrow \infty$ in the space $Y_{(\nu')}$. Hence it follows that $B \subset S - S$, where S is compact in (X, τ) and $S \in \beta_c^k$.

Choose $\delta_0 > 0$ such that $x + \delta h \in U$, $x + \delta u \in U$, and $x + \delta v \in U$, where $|\delta| \leq \delta_0$ and $h = u - v$, $h \in B$, $u, v \in S$. We introduce the notation

$$\omega(x, \delta h) = A(x + \delta h) - A(x) - A'(x)\delta h.$$

It is obvious that

$$\omega(x, \delta h) = A(x + \delta h) - A(x + \delta h + \delta v) + A(x + \delta h + \delta v) - A(x) - A'(x)\delta h.$$

Hence by the continuity of $A'(x)$ at the point x , we obtain the following identities:

$$\begin{aligned} \omega(x, \delta h) &= -\int_0^1 A'(x + \delta h + t\delta v)\delta v dt + \int_0^1 A'(x + t(\delta h + \delta v))(\delta h + \delta v) dt \\ &\quad - \int_0^1 A'(x)\delta h dt \\ &= \int_0^1 [A'(x) - A'[x + \delta h + t\delta v]]\delta v dt \\ &\quad + \int_0^1 [A'(x + t\delta u) - A'(x)]\delta u dt. \end{aligned} \tag{4}$$

Again, by the continuity of $A'(x)$, there exists a neighbourhood of zero P in the space (X, τ) such that for all $u, v \in S$, we have the inclusions

$$[A'(x) - A'(x + P)]v \subset \frac{1}{2}W,$$

and

$$[A'(x + P) - A'(x)]u \subset \frac{1}{2}W.$$

Since the set S is bounded in (X, τ) , there exists $\delta_W > 0$ such that for $|\delta| < \delta_W$, we have (as a result of (4)) the inclusions

$$\frac{\omega(x, \delta h)}{\delta} \subset \frac{1}{2}W + \frac{1}{2}W = W.$$

Now (3) follows from these inclusions. The theorem is proved.

Corollary 2. *Let (X, τ) be the strict inductive limit of the sequence $\{X_n : n = 1, 2, \dots\}$ of Fréchet-Montel spaces and let K be a generating closed cone in (X, τ) . Then if the weak derivative $A'(x)$ with respect to the cone K of the operator $A : X \rightarrow Y$ is a continuous mapping into $L_\beta(X, Y)$ on the open neighbourhood U of the point x , it is the bounded derivative of the operator A at the points $x \in U$.*

3. The Lagrange Function in *SH*-Spaces

In this section, we give the application already mentioned of Theorem 1 to the linear optimization problem in an LCS. Suppose that it is required to minimize the functional $f(x)$ under the condition $Ax \geq y_0$, where X and Y are LCSs, $A : X \rightarrow Y$ is a continuous linear operator, f is a continuous linear functional on X ; (inequalities in Y are to be understood in the sense of the ordering defined by the cone K).

We recall [5] that a point $(x_0, y'_0) \in X \times K_{Y'}$ is called a *saddle point* of the Lagrange function

$$H(x, y') = f(x) - y'(Ax - y_0),$$

if

$$H(x, y'_0) \geq H(x_0, y'_0) \geq H(x_0, y') \quad (x \in X, y' \in K_{Y'}).$$

Below we denote by Y_0 the linear hull in Y of an element y_0 and the subspace AX and by M the set $\{x : Ax \geq y_0\}$.

Theorem 3. *Let Y be a sequentially complete bornological SH -space and let K_Y be a generating closed cone in Y ; suppose moreover that $Y = Y_0 - K_Y$. Then, the functional f attains a minimum on the set M if and only if the corresponding Lagrange function $H(x, y')$ has a saddle point (x_0, y'_0) ($x \in X, y' \in K_{Y'}$).*

For the proof, it is enough to refer to Corollary 1 and Theorem 9 of [5].

4. Conclusion

Using the closed graph theorem is the important resource for applying of space Y . Such condition for space Y is being SH -space of Smirnov. In particular, this class contains of Fréchet spaces and spaces $D'(\mathbb{R}^n)$ of generalized functions. So such approach lead to an expansion of mathematical models for economic tasks of optimum control in locally convex spaces.

References

- [1] V. I. Averbukh and O. G. Smolyanov, Theory of differentiation in linear topological spaces (Russian), Uspehi. Mat. Nauk. 22 (1967), 201-258.
- [2] I. F. Danilenko, On nonoblate cones in locally convex spaces (Russian), Vestnik Leningrad Univ. 7 (1972), 29-35.
- [3] M. De Wilde, Réseaux dans les espaces linéaires à semi-normes, Mém. Soc. Royale Sci. Liège. 18 (1969), 1-144.
- [4] R. Edwards, Functional Analysis: Theory and Applications (Russian), Mir, Moscow, 1971 (transl. from English Edition: New York, Holt, Rinehart and Winston, 1965).
- [5] V. L. Levin, Conditions for an extremum in infinite-dimensional linear problems with operator restrictions (Russian), Issled. Mat. Programm., Novosibirsk (1972), 159-197.
- [6] E. I. Smirnov, Hausdorff Spectra in Functional Analysis, Springer-Verlag, London, 2002.
- [7] E. I. Smirnov, On the Hausdorff limit of locally convex spaces (Russian), Sibirsk. Mat. Zh. 28 (1987), 153.
- [8] E. I. Smirnov, Using homological methods on the base of iterated spectra in functional analysis, Vladikavkaz Mathematical Journal 14(4) (2012), 73-82.

